Comparison of GMI output for lightning simulations with SHADOZ O₃ profiles


Dale Allen and Ken Pickering

- For year 1 of the GMI simulations, hourly model output was archived on 20 pressure levels at the location of SHADOZ O₃ sondes
- Ascension Island and Nairobi, Kenya (1998-2004 data; one sonde per week with gaps)
- All model output and sonde data were used to create:
- 1. Vertical profiles (Model versus SHADOZ for no lightning, default lightning, and new lightning simulations)
- 2. Curtain (time series) plots
- 3. Probability distribution functions
- 4. Percent lightning contribution plots




Anne. M. Thompson, PI


site: Ascension (-7.98 deg lat, -14.42 deg lon), model: DAO




site: Ascension (-7.98 deg lat, -14.42 deg lon), model: FVGCM





site: Ascension (-7.98 deg lat, -14.42 deg lon), lightning flag: v2L2





site: Ascension (-7.98 deg lat, -14.42 deg lon), lightning flag: v2L2















03 PDF at ASC(-7.98N - 14.42E): 350.000 hPa













03 PDF at ASC(-7.98N - 14.42E): 350.000 hPa














03 PDF at ASC(-7.98N - 14.42E): 350.000 hPa













Summary of Ascension Island comparison

- DAO:
- In upper troposphere, 5-15 ppbv more O₃ in new run during Sep-Nov, bringing model closer to observed values
- Lightning contribution of 35-45% to O₃ in upper troposphere
- GISS:
- Upper tropospheric O₃ in new lightning run 5-15 ppbv < O₃ in default run for November – June period; GISS cloud tops too low
- Lightning contribution of 35-50% to O₃ in upper troposphere
- FVGCM:
- With default or new lightning- High-bias above ~300 hPa.
- (unobserved) linear increase with height during several months
- Lightning contribution of 40 to >60% to O₃ in upper troposphere
- ALL:
- Improvement in shape of frequency distribution of 350 hPa ozone mixing ratios


site: Ascension (-7.98 deg lat, -14.42 deg lon), model: DAO




site: Ascension (-7.98 deg lat, -14.42 deg lon), model: GISS




site: Ascension (-7.98 deg lat, -14.42 deg lon), model: GISS




site: Ascension (-7.98 deg lat, -14.42 deg lon), model: FVGCM




site: Ascension (-7.98 deg lat, -14.42 deg lon), model: FVGCM




site: Nairobí (-01.27 deg lat, +36.80 deg lon), model: DAO




site: Nairobí (-01.27 deg lat, +36.80 deg lon), model: DAO




site: Nairobí (-01.27 deg lat, +36.80 deg lon), model: GISS




site: Nairobí (-01.27 deg lat, +36.80 deg lon), model: GISS




site: Nairobi (-01.27 deg lat, +36.80 deg lon), model: FVGCM




site: Nairobi (-01.27 deg lat, +36.80 deg lon), model: FVGCM





site: Nairobí (-01.27 deg lat, +36.80 deg lon), lightning flag: v2L2





site: Nairobi (-01.27 deg lat, +36.80 deg lon), lightning flag: v2L2















03 PDF at NAI(-01.27N + 36.80E); 350.000 hPa













03 PDF at NAI(-01.27N + 36.80E): 350.000 hPa









03 PDF at NAI(-01.27N + 36.80E); 350.000 hPa

